



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

conglomerate, sandstone, shales, and "marls") and "Leroux formation" (shales, with some sandstone and calcareous beds); and Quaternary moraines and alluvium. The thickness of this generalized section is about 2,500 feet. An unconformity occurs between the Kaibab and the Moencopic, between the latter and the "Lithodendron formation," and between the Triassic and later rocks. The Moencopic formation is a fluvatile or shallow-water deposit; the Triassic beds are continental deposits. The major structural feature of the region is a very flat anticline which trends N. 30° W.

Chap. iii gives detailed descriptions of the volcanoes and lava fields. Three general periods of volcanic activity are recognized: (1) widespread basaltic eruptions from small cones, (2) eruptions of lavas (andesites to rhyolites) to form a few large cones, and laccolithic intrusions, (3) extrusion of basalt (less widespread but more cones built up than in the first named). San Francisco Mountain, which is the principal feature of the area, is composed of "lavas and breccias belonging to five distinct stages of eruption."

"The Geologic History of the Volcanic Field and Adjacent Country" is given in chap. iv. The volcanic activity of the first period occurred in the late Pliocene after the peneplanation of the region, that of the second period took place in the early Quaternary during or after the mature dissection of the area, and that of the third period during the latter part of the Quaternary subsequent to broad regional uplift. There was folding and flexing during the latter half or at the close of the Eocene. Faulting occurred at the close of the Miocene, at the close of the Pliocene, and during the middle or latter part of the Quaternary.

The last two chapters, v and vi, are devoted respectively to petrography and petrology.

V. O. T.

---

*Transactions of the American Institute of Mining Engineers.* Vol. L.  
New York, 1915. Pp. 1008.

Material for this volume was presented at the Pittsburgh meeting in October, 1914. Three topics include the major portion of the volume: (1) iron, geology, and metallurgy; (2) coal and coke with by-products; and (3) petroleum. The volume contains less purely geological matter than either of the preceding volumes for the year. Fifty-two papers and discussions, many of which are illustrated, are included.

A. D. B.